

Kent and Medway

<u>Prescribing of Adhesive Remover Sprays/Wipes for Stoma Care</u> <u>Position Statement</u>

Kent & Medway ICB recommends:

Adhesive Remover Sprays

1st Line: StoCare® Remove Medical Adhesive Remover Spray (Can)

2nd line: Clinipeel [®] Stoma Adhesive Remover Spray (Can)

Adhesive Remover Wipes (only to be used if dexterity issues)

1st Line: Stocare® Remover Wipes

Monthly Quantity to Prescribe

Ileostomy/Urostomy: 1-2 spray cans or 30 wipes (i.e. a maximum of two

spray cans or one pack of wipes)

Colostomy: 2-4 spray cans or 30-60 wipes (i.e. a maximum of four spray cans or two packs of wipes) per 60 pouches 3-6 spray cans or 60-90 wipes per 90 pouches (i.e. a maximum of 6 spray cans or 3 packs of

wipes)

Recommendations

- For all patients, StoCare® Remove Medical Adhesive Remover Spray Can should be prescribed as first line option.
- Clinipeel ® Stoma Adhesive Remover Spray Can should be prescribed as second line option.
- Wipes should be reserved for people who lack the manual strength or dexterity to use a spray.
- Either a spray or wipes should be prescribed, not both.
- If unsure of an additional request, issue one remover spray or one box of wipes and refer patient to the stoma care nurse.
- Excessive ordering of adhesive removers should be questioned. It may signify that the person is having problems and needs a stoma assessment.

Adhesive Remover Sprays

1 st Line	Code	Price	Size
StoCare® Remove Medical Adhesive Remover Spray Can	STC100	£4.95*	50ml

2 nd Line	Code	Price	Size
Clinipeel Mint adhesive remover (Clinimed Ltd)	4502	£6.98*	50ml

Adhesive Remover Wipes (reserved for people who lack the manual strength or dexterity to use a spray)

1 st Line	Code	Price	Size
StoCare® Remove 'Non Sting' Medical Adhesive Remover	STC200	£8.50*	30 wipes
Wipes			

2 nd Line	Code	Price	Size
Lifteez Non Sting adhesive remover wipes	60321	£8.50*	30 wipes



Rationale

- Adhesive remover sprays are first line formulary choice as they are more cost-effective than wipes.
- Ileostomy/urostomy bags may be changed daily, on alternate days or every third day.
- Colostomy bags may be changed 1-3 times a day

Background

- Adhesive removers are mainly prescribed to ease discomfort and prevent skin damage when removing the stoma appliance from the skin.
- Modern aerosol sprays contain a 'bag on valve' system which ensures as much product as possible is utilised, there is less wastage, and the can will spray from different angles.

Stoma Care Nurse Team Contact Details:

		1
East Kent Stoma Care Office	Telephone: 01233 616646	Email:
		ekh-tr.stomasupport@nhs.net
Medway & Swale Stoma Office	Telephone: 01634 975291	Email:
wicdway & Swale Stoffia Office		medwayft.stomanurses@nhs.ne
Maidstone & Tunbridge Wells Stoma Office	Telephone: 01892632323 Tunbridge	Email:
		mtw-tr.stomacnsteam@nhs.net
Dartford & Gravesham Stoma Office	Telephone: 01322 428100 ext 8069	Email:
		dgn-tr.stomacare@nhs.net

References:

- 1. PrescQIPP Stoma and Continence: Adhesive Removers 2.0 (March 2018)
- 2. London Procurement Partnership; Formulary for Prescribing Adhesive Removers in Stoma Care (January 2019)

Version	1.0
Developed by	Medicines Optimisation Team, Kent & Medway Integrated Care Board Stoma Care Clinical Nurse Specialists, East Kent, Medway & Swale, Maidstone & Tunbridge Wells, Dartford & Gravesham
Date ratified	V1 August 2024 (Medicines Optimisation xxxxxx),
Review date	The recommendation is based upon the evidence available at the time of publication. This recommendation will be reviewed upon request in the light of new evidence becoming available.

^{*}Prices correct as of (01/10/24) taken from Drug Tarriff Sept 2024